

# Approximate Controllability of Fractional Stochastic Perturbed Control Systems Driven by Mixed Fractional Brownian Motion

Salah H. Abid\*, Sameer Q. Hasan, Uday J. Quaez

Mathematics department, Education College, Al-Mustansiriya University, Baghdad, Iraq  
 \*Corresponding author: abidsalah@gmail.com

Received July 07, 2015; Revised July 31, 2015; Accepted August 13, 2015

**Abstract** In this paper, the approximate controllability of nonlinear Fractional order  $0 < \alpha < 1$  Riemann-Liouville type stochastic perturbed control systems driven by mixed fractional Brownian motion in a real separable Hilbert spaces has been studied by using Krasnoselskii's fixed point theorem, stochastic analysis theory, fractional calculus and some sufficient conditions.

**Keywords:** approximate controllability, mixed fractional brownian motion, fixed point theorem, perturbed control systems, mild solution, control function

**Cite This Article:** Salah H. Abid, Sameer Q. Hasan, and Uday J. Quaez, "Approximate Controllability of Fractional Stochastic Perturbed Control Systems Driven by Mixed Fractional Brownian Motion." *American Journal of Applied Mathematics and Statistics*, vol. 3, no. 4 (2015): 168-176. doi: 10.12691/ajams-3-4-7.

## 1. Introduction

The aim main of this paper is to study the principle concepts of the approximate controllability for complicated classes of fractional order  $0 < \alpha < 1$  Riemann-Liouville type stochastic perturbed control systems driven by mixed fractional Brownian motion The following form is the system under our consideration,

$$\begin{aligned} {}^L D_t^\alpha x(t) &= -(A + \Delta A)x(t) + Bu(t) \\ &+ F\left(t, x(t), \int_0^t h(t, s, x(s)) ds\right) \\ &+ G\left(t, x(t), \int_0^t g(s, x(s)) dW(s)\right) \frac{dW(t)}{dt} \\ &+ \sigma(t) \frac{dW^H(t)}{dt}, \quad t \in [0, T], \quad 0 < \alpha < 1 \\ {}^L D_t^{\alpha-1} x(t)|_{t=0} &= x_0 \end{aligned}$$

where,  ${}^L D_t^\alpha$  the Riemann-Liouville fractional derivative of order  $0 < \alpha < 1$ .  $-A$  is the infinitesimal generator of a compact analytic semigroup of uniformly bounded linear operators  $S(t), t \geq 0$ ,  $-\Delta A$  is a bounded linear operator in a real separable Hilbert space  $X$ . Assume that  $0 \in \rho(A + \Delta A)$ . The space  $X_\beta$  is a Hilbert such that  $X_\beta = D(A + \Delta A)^\beta$ ,  $0 \leq \beta < 1$ , equipped with norm  $\|x\|_\beta = \|(A + \Delta A)^\beta x\|_X$ .

$$\begin{aligned} x(t) &\in C_{1-\alpha}\left([0, T]; L^2(\Omega, X_\beta)\right) \\ &= \left\{x(t); t^{1-\alpha} x(t) \in C\left([0, T]; L^2(\Omega, X_\beta)\right)\right\} \end{aligned}$$

is a continuous at  $t \in [0, T]$  } with the norm  $\|x\|_{C_{1-\alpha}} = \left(\sup_{t \in [0, T]} (t^{1-\alpha} E\|x(t)\|_\beta^2)\right)^{\frac{1}{2}}$ .  $x_0$  is  $F_0$ -measurable  $X$ - valued random variable independent of  $W$  and  $W^H$  which defined on a complete probability space  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$ . The control function  $u(\cdot) \in L^2_{\mathcal{F}}([0, T]; U)$ ,  $U$  is a Hilbert space and the operator  $B$  from  $U$  into  $X$  is a bounded linear operator such that there exists constant  $L_B > 0$ ,  $\|Bu\| \leq L_B \|u\|$ .  $W = \{W(t), t \in [0, T]\}$  is a standard cylindrical Brownian motion defined on  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$  with values in a Hilbert space  $K$ . Let  $Q$  be a positive, self-adjoint and trace class operator on  $K$  and let  $L_2(K, X)$  be the space of all  $Q$ -Hilbert-Schmidt operators acting between  $K$  and  $X$  equipped with the Hilbert-Schmidt norm  $\|\cdot\|_{L_2}$ .  $W^H = \{W^H(t), t \in [0, T]\}$  is a  $Q$ -fractional Brownian motion with Hurst index  $H \in (\frac{1}{2}, 1)$  defined in a complete probability space  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, P)$  with values in a Hilbert space  $Y$ , such that  $Q$  is a positive, self-adjoint and trace class operator on  $Y$  and let  $L_2^0(Y; X)$  be the space of all  $Q$ -Hilbert-Schmidt operators acting between  $Y$  and  $X$  equipped with the Hilbert-Schmidt norm  $\|\cdot\|_{L_2^0}$ . The functions  $F: [0, T] \times X_\beta \times X_\beta \rightarrow X$ ,  $h: [0, T] \times [0, T] \times X_\beta \rightarrow X_\beta$ ,  $G: [0, T] \times X_\beta \times X_\beta \rightarrow L_2(K; X)$  and  $\sigma: [0, T] \rightarrow L_2^0(Y, X)$  are continuous functions.

Approximate controllability of stochastic control system driven by fractional Brownian motion has been interested by many authors; Sakthivel [19] study for the approximate controllability of impulsive stochastic systems with fractional Brownian motion. Guendouzi and Idrissi, [7] established and discussed the approximate controllability result of a class of dynamic control systems described by nonlinear fractional stochastic functional

differential equations in Hilbert space driven by fractional Brownian motion with Hurst parameter  $H > \frac{1}{2}$ . Ahmed [2] investigate the approximate controllability problem for the class of impulsive neutral stochastic functional differential equations with finite delay and fractional Brownian motion with Hurst parameter  $H > \frac{1}{2}$  in a Hilbert space. Abid, Hasan and Quaez [1] studied the Approximate controllability of fractional stochastic integro-differential equations which is derived by mixed type of fractional Brownian motion with Hurst parameter  $H > \frac{1}{2}$  and wiener process in real separable Hilbert space.

In this paper we will study the approximate controllability of nonlinear stochastic system. More precisely, we shall formulate and prove sufficient conditions for the Approximate controllability of Fractional order  $0 < \alpha < 1$  Riemann-Liouville type stochastic perturbed control systems driven by mixed fractional Brownian motion in a real separable Hilbert spaces.

The rest of this paper is organized as follows, in section 2, we will introduced some concepts, definitions and some lemmas of semigroup theory and fractional stochastic calculus which are useful for us here. In section 3, we will prove our main result.

## 2. Preliminaries

In this section, we introduce some notations and preliminary results, which we needed to establish our results.

### Definition (2.1), [5]:

Let  $H$  be a constant belonging to  $(0, 1)$ . A one dimensional fractional Brownian motion  $B^H = \{B^H_t, t \geq 0\}$  of Hurst index  $H$  is a continuous and centered Gaussian process with covariance function

$$E(B^H_t B^H_s) = \frac{1}{2} (t^{2H} + s^{2H} - |t - s|^{2H}), \text{ for } t, s \geq 0. (1)$$

- If  $H = \frac{1}{2}$ , then the increments of  $B^H$  are non-correlated, and consequently independent. So  $B^H$  is a Wiener Process which we denote further by  $B$ .
- If  $H \in (\frac{1}{2}, 1)$  then the increments are positively correlated.
- If  $H \in (0, \frac{1}{2})$  then the increments are negative correlated.

$B^H$  has the integral representation

$$B^H_t = \int_0^t K_H(t, s) dB(s) \tag{2}$$

where,  $B$  is a wiener process and the kernel  $K_H(t, s)$  defined as

$$K_H(t, s) = cH s^{\frac{1}{2}-H} \int_s^t (u-s)^{H-\frac{3}{2}} u^{H-\frac{1}{2}} du \tag{3}$$

$$\frac{\partial K}{\partial t}(t, s) = cH \left(\frac{t}{s}\right)^{H-\frac{1}{2}} (t-s)^{H-\frac{3}{2}} \tag{4}$$

$$cH = \left[ \frac{H(2H-1)}{\beta(2-2H, H-\frac{1}{2})} \right]^{\frac{1}{2}}, t > s \text{ and } \beta \text{ is a beta function.}$$

In the case  $H = \frac{1}{2}$ , we shall use Ito Isometry theorem

### Lemma (2.1), "Ito isometry theorem", [11]:

Let  $V [0, T]$  be the class of functions such that  $f: [0, T] \times \Omega \rightarrow \mathbb{R}$ ,  $f$  is measurable,  $\mathcal{F}_t$ - adapted and  $E \left[ \int_0^T (f(t, \omega))^2 dt \right] \leq \infty$ . Then for every  $f \in V [0, T]$ , we have

$$E \left[ \int_0^T f(t, \omega) dB(t) \right]^2 = E \left[ \int_0^T (f(t, \omega))^2 dt \right] \tag{5}$$

where  $B$  is a wiener process.

Now, we denote by  $\mathcal{E}$  the set of step functions on  $[0, T]$ . If  $\Phi \in \mathcal{E}$  then, we can write it the form as:

$$\Phi(t) = \sum_{k=1}^n a_k 1_{[t_k, t_{k+1}]}(t), \text{ where } t \in [0, T].$$

The integral of a step function  $\Phi \in \mathcal{E}$  with respect to one dimensional fractional Brownian motion is defined

$$\int_0^T \Phi(t) dB_t^H = \sum_{k=1}^n a_k (B_{t_{k+1}}^H - B_{t_k}^H),$$

where  $a_k \in \mathbb{R}$ ,

$$0 = t_1 < t_2 < \dots < t_{n+1} = T.$$

Let  $\mathcal{H}$  be the Hilbert space defined as the closure of  $\mathcal{E}$  with respect to the scalar product  $\langle 1_{[0,t]}, 1_{[0,s]} \rangle = R_H(t, s) = E(B_t^H B_s^H)$ . The mapping  $1_{[0,t]} \rightarrow \{B^H(t), t \in [0, T]\}$  can be extended to an isometry between  $\mathcal{H}$  and  $\text{span}^{L^2(\Omega)} \{B^H_t, t \in [0, T]\}$  i.e. the mapping  $\mathcal{H} \rightarrow L^2(\Omega, \mathcal{F}, P)$ ,  $\Phi \rightarrow \int_0^T \Phi(t) dB_t^H$  is isometry.

### Remark (2.1):

- If  $H = \frac{1}{2}$  and  $\mathcal{H} = L^2([0, T])$  then by use Ito isometry, we have

$$E \left( \int_0^T \Phi(t) dB \right)^2 = \int_0^T (\Phi(t))^2 dt \tag{6}$$

- If  $H > \frac{1}{2}$ , we have

$$R_H(s, t) = \frac{1}{2} (|t|^{2H} + |s|^{2H} - |t-s|^{2H}), t, s \geq 0 \tag{7}$$

$$\frac{\partial R_H}{\partial t} = H(|t|^{2H-1} - |t-s|^{2H-1}) \tag{8}$$

$$\partial R_H^2 = H(2H-1)|t-s|^{2H-2} ds dt$$

### Lemma (2.2), [6]:

For any functions  $\Phi, \varphi \in L^2[0, T] \cap L^1[0, T]$ , we have

$$i) E \left( \int_0^T \Phi(t) dB_t^H \int_0^T \varphi(s) dB_s^H \right) = H(2H-1) \int_0^T \int_0^T \Phi(t) \varphi(s) |t-s|^{2H-2} ds dt$$

$$ii) E(dB_t^H dB_s^H) = \frac{\partial R_H^2}{\partial s dt} = H(2H-1)|t-s|^{2H-2} ds dt$$

From this Lemma above, we obtain

$$E \left( \int_0^T \Phi(t) dB_t^H \right)^2 \tag{9}$$

$$= H(2H-1) \int_0^T \int_0^T \Phi(s) \Phi(t) |t-s|^{2H-2} ds dt$$

### Remark (2.2), [6]:

The space  $\mathcal{H}$  contains the set of functions  $\Phi \in L^2[0, T]$ , such that,  $\int_0^T \int_0^T \Phi(s) \Phi(t) |t-s|^{2H-2} ds dt < \infty$ , which includes  $L^{\frac{1}{H}}([0, T])$ .

Now,

Let  $\mathcal{H}$  be the Banach space of measurable functions on  $[0, T]$ , such that

$$\|\Phi\|_{\mathcal{H}}^2 = H(2H-1) \int_0^T \int_0^T \Phi(s) \Phi(t) |t-s|^{2H-2} ds dt < \infty \tag{10}$$

### Lemma (2.3), [10]:

$$L([0, T]) \subset L^{\frac{1}{H}}([0, T]) \subset \mathcal{H} \subset \mathcal{K}$$

Suppose that there exists a complete orthonormal system  $\{e_n\}_{n=1}^{\infty}$  in  $Y$ . Let  $Q \in L(Y, Y)$  be the operator defined by  $Qe_n = \lambda_n e_n$ , where  $\lambda_n \geq 0$  ( $n=1, 2, \dots$ ) are non-negative real numbers with finite trace  $\text{Tr } Q = \sum_{n=1}^{\infty} \lambda_n < \infty$ . The infinite dimensional fractional Brownian motion on  $Y$  can be defined by using covariance operator  $Q$  as

$$W_{(t)}^H = W_{Q(t)}^H = \sum_{n=1}^{\infty} \sqrt{\lambda_n} e_n B_n^H(t),$$

where  $B_n^H(t)$  are one dimensional fractional Brownian motions mutually independent on  $(\Omega, \mathcal{F}, P)$ .

In order to defined stochastic integral with respect to the  $Q$ -fractional Brownian motion. We introduce the space  $L_2^0(Y, X)$  of all  $Q$ -Hilbert- Schmidt operators that is with the inner product  $\langle \Phi, \varphi \rangle_{L_2^0} = \sum_{n=1}^{\infty} \langle \Phi e_n, \varphi e_n \rangle$  is a separable Hilbert space.

**Lemma (2.4), [10]:**

Let  $\{\Phi(t)\}_{t \in [0, T]}$  be a deterministic function with values in  $L_2^0(Y, X)$  The stochastic integral of  $\Phi$  with respect to  $W^H$  is defined by

$$\begin{aligned} \int_0^t \Phi(s) dW_{(s)}^H &= \sum_{n=1}^{\infty} \int_0^t \sqrt{\lambda_n} \Phi(s) e_n dB_{n(s)}^H \\ &= \sum_{n=1}^{\infty} \int_0^t \sqrt{\lambda_n} (\mathcal{K}_H^*(\Phi e_n))(s) dB_{n(s)} \end{aligned} \quad (11)$$

**Lemma (2.5), [10]:**

If  $\varphi: [0, b] \rightarrow L_2^0(Y, X)$  satisfies  $\int_0^T \|\varphi(s)\|_{L_2^0}^2 ds < \infty$  then the above sum in (11) is well defined as an  $X$ -valued random variable and we have

$$E \left\| \int_0^t \varphi(s) dW_{(s)}^H \right\|^2 \leq 2H t^{2H-1} \int_0^t \|\varphi(s)\|_{L_2^0}^2 ds \quad (12)$$

**Definition (2.2), [18]:**

The Riemann - Liouville derivative of order  $\alpha > 0$  with lower limit zero for a function  $f$  can be written as:

$${}^L D^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \frac{d^n}{dt^n} \int_0^t \frac{f(s)}{(t-s)^{\alpha+1-n}} ds \quad (13)$$

where,  $t > 0, n-1 < \alpha < n$ .

**Definition (2.4), [18]:**

The Laplace transform of the Riemann-Liouville fractional derivation of order  $\alpha > 0$  gives as:

$$\mathcal{L}\{L D_t^\alpha f(t)\} = \lambda^\alpha \mathcal{L}\{f(t)\}(\lambda) - \sum_{k=0}^{n-1} \lambda^k [D_t^{\alpha-k-1} f(t)]_{t=0}(\lambda) \quad (14)$$

where,  $n-1 < \alpha < n$ .

**Lemma (2.6), [17]:**

Let  $-A$  be the infinitesimal generator of an analytic semigroup  $S(t), t \geq 0$  on a Hilbert space  $X$ . If  $-\Delta A$  is a bounded linear operator on  $X$  then  $-(A+\Delta A)$  is the infinitesimal generator of an analytic semigroup  $\check{T}(t), t \geq 0$  on  $X$ .

**Remark (2.3):**

Assume that  $\check{T}(t), t \geq 0$  is a compact analytic semigroup of uniformly bounded operators in  $X$ , that is, there exists  $M > 1$  such that  $\|\check{T}(t)x\| \leq M$ .

**Definition (2.5):**

An  $X_\beta$ -valued process  $x(t)$  is called a mild solution of the system (1) if  $x(t) \in C_{1-\alpha}([0, T]; L^2(\Omega, X_\beta))$  and, for  $t \in [0, T]$  satisfies the integral equation

$$\begin{aligned} x(t) &= \check{T}_\alpha(t)(t)^{\alpha-1} x_0 \\ &+ \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} B u(s) ds \\ &+ \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} F \left( s, x(s), \int_0^s h(s, r, x(r)) dr \right) ds \\ &+ \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} G \left( s, x(s), \int_0^s g(s, x(r)) dW_{(r)} \right) dW_{(s)} \\ &+ \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \sigma(s) dW_{(s)}^H \end{aligned} \quad (15)$$

where,  $\check{T}_\alpha(t) = \int_0^\infty \alpha r M_\alpha(r) T(t^\alpha r) dr$   
 $M_\alpha(r)$  is a Mainardi's function.

**Lemma (2.7):**

If  $\check{T}(t), t \geq 0$  is a compact analytic semigroup then the family of operators  $\check{T}_\alpha(t), t \geq 0$  have the following properties:

- i. For any fixed  $t \geq 0$ , the operator  $\check{T}_\alpha(t)$  is a linear and bounded, i.e. for any  $x \in X$ , there exists  $M \geq 1$  such that  $\|\check{T}_\alpha(t)x\|_X \leq \frac{M}{\Gamma(\alpha)} \|x\|_X$ .
- ii. For any  $x \in X_\beta$ , there exists  $M \geq 1$  such that  $\|\check{T}_\alpha(t)x\|_\beta \leq \frac{M}{\Gamma(\alpha)} \|x\|_\beta$ .
- iii. For any  $x \in X, \|\check{T}_\alpha(t)x\|_\beta \leq \check{N}_{\alpha, \beta} t^{-\alpha\beta} \|x\|_X$  where,  $\check{N}_{\alpha, \beta} = \alpha \check{K}_\beta \frac{\Gamma(2-\beta)}{\Gamma(\alpha(1-\beta)+1)}$
- iv.  $\{\check{T}_\alpha(t), t \geq 0\}$  is a strongly continuous, which mean that for every  $x \in X$  and  $0 \leq t_1 < t_2 \leq T$  then  $\|\check{T}_\alpha(t_2)x - \check{T}_\alpha(t_1)x\|_X \rightarrow 0$  if  $t_2 \rightarrow t_1$ .
- v. The operator  $\check{T}_\alpha(t)$  is a compact operator in  $X$  for  $t > 0$ .

### 3. Main Result of the Approximately Controllable

In this section, we formulate and prove the result on approximate controllability of nonlinear fractional stochastic perturbed control system driven by mixed fractional Brownian motion in (1). To establish our results, we introduce the following assumptions:

- a) The operator  $\check{T}(t)$  is a compact for any  $t > 0$ .
- b) The linear fractional order system of corresponding the system (3.41) which has following form:

$$\begin{aligned} {}^L D_t^\alpha x(t) &= Ax(t) + Bu(t), t \in [0, T] \\ {}^L D_t^{\alpha-1} x(t)|_{t=0} &= x_0, \frac{1}{2} < \alpha < 1 \end{aligned} \quad (16)$$

is an approximately controllable on  $[0, T]$ .

- c) The functions  $F: [0, T] \times X_\beta \times X_\beta \rightarrow X, h: [0, T] \times [0, T] \times X_\beta \rightarrow X_\beta$  are satisfying linear growth and Lipschitz conditions. This mean that, for any  $x, y \in X_\beta$ , there exists positive constants  $K_1, K_2 > 0$  and  $K_3, K_4 > 0$  such that

$$\begin{aligned} &\left\| F \left( t, x(t), \int_0^t h(t, s, x(s)) ds \right) - F \left( t, y(t), \int_0^t h(t, s, y(s)) ds \right) \right\|_X \leq K_1 \|x - y\|_\beta \\ &\left\| F \left( t, x(t), \int_0^t h(t, s, x(s)) ds \right) \right\|_X^2 \leq K_2 (1 + \|x\|_\beta^2) \\ &\|h(t, s, x) - h(t, s, y)\|_\beta^2 \leq K_3 \|x - y\|_\beta^2 \\ &\|h(t, s, x)\|_\beta^2 \leq K_4 (1 + \|x\|_\beta^2) \end{aligned}$$

Also, F is a uniformly bounded. In other word, there exists  $D_1 > 0$  such that

$$\left\| F\left(t, x(t), \int_0^t h(t, s, x(s)) ds\right) \right\|_X^2 < D_1, \text{ for } t \in [0, T]$$

d) The function  $\sigma: [0, T] \rightarrow L^0_2(Y; X)$  satisfies, for every  $t \in [0, T]$

$\int_0^t \|\sigma(s)\|_{L^0_2}^2 ds < \infty$  and there exists  $C_1 > 0$  such that  $\sup_{t \in [0, T]} \|\sigma(s)(t)\|_{L^0_2}^2 \leq C_1$ .

e)  $\tilde{G}$  is  $F_t$  - adapted with respect to  $t$ , such that, for every  $t \in [0, T]$ , satisfy the following:

- i.  $E \left\| (\tilde{G}x)(s) \right\|_{L^2}^2$  Exists.
- ii.  $\int_0^t E \left\| (\tilde{G}x)(t) \right\|_{L^2}^2 ds < \infty$ .
- iii. There exists  $D_2 > 0$ , such that

$$\sup_{t \in [0, T]} E \left\| (\tilde{G}x)(t) \right\|_{L^2}^2 < D_2.$$

where,  $(\tilde{G}x)(t) = G\left(t, x(t), \int_0^t g(s, x(s)) dW_{(s)}\right)$

**Definition (3.1):**

The system (1) is said to be approximately controllable on  $[0, T]$  if the reachable set  $\mathfrak{R}(T)$  is dense in the space  $L^2(\Omega, X)$ . This mean that  $\overline{\mathfrak{R}(T)} = L^2(\Omega, X)$ . where,  $\mathfrak{R}(T) = \{x(T, u) : u \in L^2_{\mathfrak{F}}([0, T]; U)\}$

Now,

The controllability operator  $\Gamma_s^T$  associated with control system (16) is defined by

$$\Gamma_s^T = \int_s^T (T-t)^{\alpha-1} \check{T}_\alpha(T-t) B B^* \check{T}_\alpha(T-t) dt \quad (17)$$

Also, for any  $\theta > 0$  and  $0 \leq s < T$ , the operator  $R(\theta, \Gamma_s^T)$  is defined by

$$R(\theta, \Gamma_s^T) = (\theta I + \Gamma_s^T)^{-1} \quad (18)$$

where,  $B^*$  and  $\check{T}_\alpha^*$  are the adjoint operators for  $B$  and  $\check{T}_\alpha$  respectively.

**Lemma (3.1), [12]:**

The linear fractional order deterministic system in (1) is an approximately controllable on  $[0, T]$  if and only if the operator  $\theta R(\theta, \Gamma_0^T) \rightarrow 0$  as  $\theta \rightarrow 0^+$ ,  $\|\theta R(\theta, \Gamma_s^T)\| \leq 1$ .

**Lemma (3.2), [13]:** For any  $x_T \in L^2(\Omega, \mathfrak{F}_T, X_\beta)$ , there exists

$\hat{H} \in L^2(\Omega; L^2([0, T]; L^0_2(Y, X)))$  and  $\phi \in L^2(\Omega; L^2([0, T]; L_2(K; X)))$ , such that

$$x_T = E x_T + \int_0^T \phi(s) dW_{(s)} + \int_0^T \hat{H}(s) dW_{(s)}^H \quad (19)$$

where,  $\sup_{t \in [0, T]} E \|\phi(t)\|_{L^2}^2 \leq \check{C}_1$ ,

$$\sup_{t \in [0, T]} E \|\hat{H}(t)\|_{L^0_2}^2 \leq \check{C}_2$$

Now, For any  $\theta > 0$  and any  $x_T \in L^2(\Omega, \mathfrak{F}_T, X_\beta)$ , we defined the control function of the system (1) in the following form:

$$\begin{aligned} u^\theta(t, x) &= B^* \check{T}_\alpha^*(T-t) R(\theta, \Gamma_0^T) \left( E x_T - \check{T}_\alpha(T)(T)^{\alpha-1} x_0 \right) \\ &+ B^* \check{T}_\alpha^*(T-t) \int_0^t R\left(\theta, \Gamma_s^T\right) \phi(s) dW_{(s)} \\ &+ B^* \check{T}_\alpha^*(T-t) \int_0^t R\left(\theta, \Gamma_s^T\right) \hat{H}(s) dW_{(s)}^H \end{aligned}$$

$$\begin{aligned} &- B^* \check{T}_\alpha^*(T-t) \int_0^t \left[ R\left(\theta, \Gamma_s^T\right) \check{T}_\alpha(T-s)(T-s)^{\alpha-1} \right. \\ &\quad \left. F\left(s, x(s), \int_0^s h(s, r, x(r)) dr\right) \right] ds \\ &- B^* \check{T}_\alpha^*(T-t) \int_0^t \left[ R\left(\theta, \Gamma_s^T\right) \check{T}_\alpha(T-s)(T-s)^{\alpha-1} \right. \\ &\quad \left. G\left(s, x(s), \int_0^s g(v, x(v)) dW_{(v)}\right) \right] dW_{(s)} \quad (20) \\ &- B^* \check{T}_\alpha^*(T-t) \int_0^t \left[ R\left(\theta, \Gamma_s^T\right) \check{T}_\alpha(T-s) \right. \\ &\quad \left. (T-s)^{\alpha-1} \sigma(s) \right] dW_{(s)}^H \end{aligned}$$

**Lemma (3.3):** There exists positive real constant  $N_C$ , such that for all  $x \in C([0, T]; L^2(\Omega, X_\beta))$ ,

$$E \|u^\theta(t, x)\|^2 \leq N_C \quad (21)$$

where,

$$\begin{aligned} N_C &= \frac{12L_B^2 M^4}{\theta^2 (\Gamma(\alpha))^4} (E \|x_T\|_\beta^2 + (T)^{2\alpha-2} E \|x_0\|_\beta^2) \\ &+ 6 \frac{L_B^2}{\theta^2 (\Gamma(\alpha))^2} M^2 T \check{C}_1 + \frac{12HT^{2H} L_B^2 M^2}{\theta^2 (\Gamma(\alpha))^2} \check{C}_2 \\ &+ \frac{6L_B^2 M^2 T^{2\alpha-2\alpha\beta}}{\theta^2 (\Gamma(\alpha))^2 (1-2\alpha\beta)(2\alpha-1)} \check{N}_{\alpha, \beta}^2 D_1 \\ &+ \frac{6L_B^2 M^2 T^{2\alpha-2\alpha\beta}}{\theta^2 (\Gamma(\alpha))^2 (1-2\alpha\beta)(2\alpha-1)} \check{N}_{\alpha, \beta}^2 D_2 \\ &+ \frac{12HL_B^2 M^2 T^{2\alpha-2\alpha\beta+2H-1}}{\theta^2 (\Gamma(\alpha))^2 (2\alpha-1)(1-2\alpha\beta)} \check{N}_{\alpha, \beta}^2 C_1 \end{aligned}$$

**Proof**

Let  $x \in C([0, T]; L^2(\Omega, X_\beta))$  and  $T > 0$  be a fixed. From the equation (21), we have:

$$\begin{aligned} &E \|u^\theta(t, x)\|^2 \\ &\leq 6E \left\| B^* \check{T}_\alpha^*(T-t) R(\theta, \Gamma_0^T) \left( E x_T - \check{T}_\alpha(T)(T)^{\alpha-1} x_0 \right) \right\|^2 \\ &+ 6E \left\| B^* \check{T}_\alpha^*(T-t) \int_0^t R\left(\theta, \Gamma_s^T\right) \phi(s) dW_{(s)} \right\|^2 \\ &+ 6E \left\| B^* \check{T}_\alpha^*(T-t) \int_0^t R\left(\theta, \Gamma_s^T\right) \hat{H}(s) dW_{(s)}^H \right\|^2 \\ &+ 6E \left\| B^* \check{T}_\alpha^*(T-t) \int_0^t \left[ R\left(\theta, \Gamma_s^T\right) \check{T}_\alpha(T-s)(T-s)^{\alpha-1} \right. \right. \\ &\quad \left. \left. F\left(s, x(s), \int_0^s h(s, r, x(r)) dr\right) \right] ds \right\|^2 \\ &+ 6E \left\| B^* \check{T}_\alpha^*(T-t) \int_0^t \left[ R\left(\theta, \Gamma_s^T\right) \check{T}_\alpha(T-s)(T-s)^{\alpha-1} \right. \right. \\ &\quad \left. \left. G\left(s, x(s), \int_0^s g(v, x(v)) dW_{(v)}\right) \right] dW_{(s)} \right\|^2 \end{aligned}$$

$$+6E \left\| B^* \check{T}_\alpha^* (T-t) \int_0^t \left[ \begin{array}{l} R(\theta, \Gamma_s^T) \check{T}_\alpha (T-s) \\ (T-s)^{\alpha-1} \sigma(s) \end{array} \right] dW_{(s)}^H \right\|^2$$

Applying Holder's inequality and by using Ito isometry, Lemmas (2.5), (2.7) and the assumptions (a)-(e), we obtain

$$E \|u^\theta(t, x)\|^2 \leq N_C$$

where,

$$\begin{aligned} N_C &= \frac{12L_B^2 M^4}{\theta^2 (\Gamma(\alpha))^4} (E \|x_T\|_\beta^2 + (T)^{2\alpha-2} E \|x_0\|_\beta^2) \\ &+ 6 \frac{L_B^2}{\theta^2 (\Gamma(\alpha))^2} M^2 T \check{C}_1 + \frac{12HT^{2H} L_B^2 M^2}{\theta^2 (\Gamma(\alpha))^2} \check{C}_2 \\ &+ \frac{6L_B^2 M^2 T^{2\alpha-2\alpha\beta}}{\theta^2 (\Gamma(\alpha))^2 (1-2\alpha\beta)(2\alpha-1)} \check{N}_{\alpha,\beta}^2 D_1 \\ &+ \frac{6L_B^2 M^2 T^{2\alpha-2\alpha\beta}}{\theta^2 (\Gamma(\alpha))^2 (1-2\alpha\beta)(2\alpha-1)} \check{N}_{\alpha,\beta}^2 D_2 \\ &+ \frac{12HL_B^2 M^2 T^{2\alpha-2\alpha\beta+2H-1}}{\theta^2 (\Gamma(\alpha))^2 (2\alpha-1)(1-2\alpha\beta)} \check{N}_{\alpha,\beta}^2 C_1 \end{aligned}$$

Now,

For any  $\theta > 0$ , consider the operator  $\Psi_\theta$  on  $C_{1-\alpha}([0, T]; L^2(\Omega, X_\beta))$  defined as follows:

$$\begin{aligned} (\Psi_\theta x)(t) &= \check{T}_\alpha(t)(t)^{\alpha-1} x_0 \\ &+ \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} Bu^\theta(t, x) ds \\ &+ \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} F\left(s, x(s), \int_0^s h(s, r, x(r)) dr\right) ds \\ &+ \int_0^t \left[ \begin{array}{l} \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \\ G\left(s, x(s), \int_0^s g(v, x(v)) dW_{(v)}\right) \end{array} \right] dW_{(s)} \\ &+ \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \sigma(s) dW_{(s)}^H \end{aligned} \tag{22}$$

Also, for any  $\delta > 0$ , the subset  $B_\delta$  of  $C_{1-\alpha}([0, T]; L^2(\Omega, X_\beta))$  is define as  $B_\delta = \{x(t) \in C_{1-\alpha}([0, T]; L^2(\Omega, X_\beta)) : \|x(t)\|_{C_{1-\alpha}}^2 \leq \delta\}$ .

**Lemma (3.4):**

for any  $\theta > 0$ , There exists  $\check{\delta} > 0$  such that  $\Psi_\theta(B_{\check{\delta}}) \subset B_{\check{\delta}}$ .

**Proof:**

To prove that there exists  $\delta > 0$  such that  $\Psi_\theta(B_\delta) \subset B_\delta$ , in other word,  $\|\Psi_\theta x(t)\|_{C_{1-\alpha}}^2 \leq \delta$ , for each  $x(t) \in B_\delta$ .

Suppose that this is not true, then for each  $\delta > 0$ , there exists  $x(t) \in B_\delta$  such that  $\|\Psi_\theta x(t)\|_{C_{1-\alpha}}^2 > \delta$ , for  $t \in [0, T]$ ,  $t$  may depending upon  $\delta$ . However, on the other hand, we have

$$t^{2-2\alpha} E \|\Psi_\theta x(t)\|_\beta^2 \leq 5t^{2-2\alpha} E \left\| \check{T}_\alpha(t)(t)^{\alpha-1} x_0 \right\|_\beta^2$$

$$\begin{aligned} &+ 5t^{2-2\alpha} E \left\| \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} Bu^\theta(s, x) ds \right\|_\beta^2 \\ &+ 5t^{2-2\alpha} E \left\| \int_0^t \left[ \begin{array}{l} \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \\ F\left(s, x(s), \int_0^s h(s, r, x(r)) dr\right) \end{array} \right] ds \right\|_\beta^2 \\ &+ 5t^{2-2\alpha} E \left\| \int_0^t \left[ \begin{array}{l} \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \\ G\left(s, x(s), \int_0^s g(v, x(v)) dW_{(v)}\right) \end{array} \right] dW_{(s)} \right\|_\beta^2 \\ &+ 5t^{2-2\alpha} E \left\| \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \sigma(s) dW_{(s)}^H \right\|_\beta^2 \end{aligned}$$

Applying Holder's inequality and by using Ito isometry, Lemmas (2.5), (2.7), (3.3) and the assumptions (a)-(e), we obtain

$$\begin{aligned} &\sup_{t \in [0, T]} t^{2-2\alpha} E \|\Psi_\theta x(t)\|_\beta^2 \\ &\leq 5 \frac{M^2}{(\Gamma(\alpha))^2} E \|x_0\|_\beta^2 + \frac{5L_B^2 T^{2-2\alpha\beta} N_C \check{N}_{\alpha,\beta}^2}{(2\alpha-1)(1-2\alpha\beta)} \\ &+ \frac{5T^{2-2\alpha\beta} D_1 \check{N}_{\alpha,\beta}^2}{|(2\alpha-1)(1-2\alpha\beta)|} + \frac{5T^{2-2\alpha\beta} \check{N}_{\alpha,\beta}^2 D_2}{|(2\alpha-1)(1-2\alpha\beta)|} \\ &+ \frac{10HT^{1+2H-2\alpha\beta}}{|(2\alpha-1)(1-2\alpha\beta)|} \check{N}_{\alpha,\beta}^2 C_1 \end{aligned}$$

Hence,

$$\begin{aligned} &\|\Psi_\theta x(t)\|_{C_{1-\alpha}}^2 \\ &\leq 5 \frac{M^2}{(\Gamma(\alpha))^2} E \|x_0\|_\beta^2 + \frac{5L_B^2 T^{2-2\alpha\beta} N_C \check{N}_{\alpha,\beta}^2}{(2\alpha-1)(1-2\alpha\beta)} \\ &+ \frac{5T^{2-2\alpha\beta} D_1 \check{N}_{\alpha,\beta}^2}{|(2\alpha-1)(1-2\alpha\beta)|} + \frac{5T^{2-2\alpha\beta} \check{N}_{\alpha,\beta}^2 D_2}{|(2\alpha-1)(1-2\alpha\beta)|} \\ &+ \frac{10HT^{1+2H-2\alpha\beta}}{|(2\alpha-1)(1-2\alpha\beta)|} \check{N}_{\alpha,\beta}^2 C_1 \end{aligned}$$

Therefore,

$$\begin{aligned} \delta &< 5 \frac{M^2}{(\Gamma(\alpha))^2} E \|x_0\|_\beta^2 + \frac{5L_B^2 T^{2-2\alpha\beta} N_C \check{N}_{\alpha,\beta}^2}{(2\alpha-1)(1-2\alpha\beta)} \\ &+ \frac{5T^{2-2\alpha\beta} D_1 \check{N}_{\alpha,\beta}^2}{|(2\alpha-1)(1-2\alpha\beta)|} + \frac{5T^{2-2\alpha\beta} \check{N}_{\alpha,\beta}^2 D_2}{|(2\alpha-1)(1-2\alpha\beta)|} \\ &+ \frac{10HT^{1+2H-2\alpha\beta}}{|(2\alpha-1)(1-2\alpha\beta)|} \check{N}_{\alpha,\beta}^2 C_1 \end{aligned}$$

By dividing both sides of above inequality by  $\delta$  and taking the limit as  $\delta \rightarrow \infty$ , which is a contradiction. Thus, for each  $\theta > 0$ , there exists positive number  $\check{\delta}$  such that  $\Psi_\theta(B_{\check{\delta}}) \subset B_{\check{\delta}}$ .

Now,

Let  $\Psi_\theta = \Psi_1 + \Psi_2$  where,

$$(\Psi_1 x)(t) = \int_0^t \left[ \check{T}_\alpha(t-s)(t-s)^{\alpha-1} F\left(s, x(s), \int_0^s h(s, r, x(r)) dr\right) \right] ds \quad (23)$$

$$\begin{aligned} (\Psi_2 x)(t) &= \check{T}_\alpha(t)(t)^{\alpha-1} x_0 \\ &+ \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} Bu^\theta(s, x) ds \\ &+ \int_0^t \left[ \check{T}_\alpha(t-s)(t-s)^{\alpha-1} G\left(s, x(s), \int_0^s g(v, x(v)) dW_{(v)}\right) \right] dW_{(s)} \\ &+ \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \sigma(s) dW_{(s)}^H \end{aligned} \quad (24)$$

**Lemma (3.5):**

Assume that the assumptions (a) – (e) hold, then for any  $\theta > 0$ , and for any  $x, y \in B_{\check{\delta}}$ ,  $(\Psi_1 y)(t) + (\Psi_2 x)(t) \in B_{\check{\delta}}$ , for  $t \in [0, T]$ .

**Proof**

Let  $x, y \in B_{\check{\delta}}$  and  $0 \leq t \leq T$ , we have

$$\begin{aligned} &t^{2-2\alpha} E \|(\Psi_1 y)(t) + (\Psi_2 x)(t)\|_\beta^2 \\ &\leq 5t^{2-2\alpha} E \left\| \check{T}_\alpha(t)(t)^{\alpha-1} x_0 \right\|_\beta^2 \\ &+ 5t^{2-2\alpha} E \left\| \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} Bu^\theta(s, x) ds \right\|_\beta^2 \\ &+ 5t^{2-2\alpha} E \left\| \int_0^t \left[ \check{T}_\alpha(t-s)(t-s)^{\alpha-1} F\left(s, y(s), \int_0^s h(s, r, y(r)) dr\right) \right] ds \right\|_\beta^2 \\ &+ 5t^{2-2\alpha} E \left\| \int_0^t \left[ \check{T}_\alpha(t-s)(t-s)^{\alpha-1} G\left(s, x(s), \int_0^s g(v, x(v)) dW_{(v)}\right) \right] dW_{(s)} \right\|_\beta^2 \\ &+ 5t^{2-2\alpha} E \left\| \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \sigma(s) dW_{(s)}^H \right\|_\beta^2 \end{aligned}$$

Applying Holder’s inequality and by using Ito isometry, Lemmas (2.5), (2.7), (3.3) and the assumptions (a)-(e), we obtain

$$\begin{aligned} &t^{2-2\alpha} E \|(\Psi_1 y)(t) + (\Psi_2 x)(t)\|_\beta^2 \\ &\leq 5 \frac{M^2}{(\Gamma(\alpha))^2} E \|x_0\|_\beta^2 + \frac{5L_B^2 T^{2-2\alpha\beta} N_C \check{N}_{\alpha, \beta}^2}{(2\alpha-1)(1-2\alpha\beta)} \\ &+ \frac{5T^{2-2\alpha\beta} D_1 \check{N}_{\alpha, \beta}^2}{|(2\alpha-1)(1-2\alpha\beta)|} + \frac{5T^{2-2\alpha\beta} \check{N}_{\alpha, \beta}^2 D_2}{|(2\alpha-1)(1-2\alpha\beta)|} \\ &+ \frac{10HT^{1+2H-2\alpha\beta}}{|(2\alpha-1)(1-2\alpha\beta)|} \check{N}_{\alpha, \beta}^2 C_1 \end{aligned}$$

which mean  $\|(\Psi_1 y)(t) + (\Psi_2 x)(t)\|_{C_{1-\alpha}}^2 \leq \check{\delta}$ , then  $(\Psi_1 y)(t) + (\Psi_2 x)(t) \in B_{\check{\delta}}$

**Lemma (3.6):**

For any  $t \in [0, T]$ , the operator  $\Psi_1$  is a contraction on  $B_{\check{\delta}}$ , provided that  $\gamma = \frac{\check{N}_{\alpha, \beta}^2 T^{2-2\alpha\beta} K_1}{|(2\alpha-1)(1-2\alpha\beta)|} < 1$

**Proof:**

Let  $t \in [0, T]$ , and  $y_1, y_2 \in B_{\check{\delta}}$ , we have

$$\begin{aligned} &t^{2-2\alpha} E \|\Psi_1 y_1(t) - \Psi_1 y_2(t)\|_\beta^2 \\ &= t^{2-2\alpha} E \int_0^t \left\{ \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \left[ F\left(s, y_1(s), \int_0^s h(s, r, y_1(r)) dr\right) - F\left(s, y_2(s), \int_0^s h(s, r, y_2(r)) dr\right) \right] \right\} ds \end{aligned}$$

By using lemma (2.7) and assumption (c), we get

$$\begin{aligned} &t^{2-2\alpha} E \|\Psi_1 y_1(t) - \Psi_1 y_2(t)\|_\beta^2 \\ &\leq \frac{t}{|2\alpha-1|} \int_0^t \check{N}_{\alpha, \beta}^2 (t-s)^{-2\alpha\beta} K_1 E \|y_1(s) - y_2(s)\|_\beta^2 ds \end{aligned}$$

By taking the supremum over  $t \in [0, T]$  for both sides, we have

$$\begin{aligned} &\sup_{t \in [0, T]} \{t^{2-2\alpha} E \|\Psi_1 y_1(t) - \Psi_1 y_2(t)\|_\beta^2\} \\ &\leq \frac{\check{N}_{\alpha, \beta}^2 T^{2-2\alpha\beta} K_1}{|(2\alpha-1)(1-2\alpha\beta)|} \sup_{t \in [0, T]} \{E \|y_1(t) - y_2(t)\|_\beta^2\} \end{aligned}$$

Therefore,

$$\begin{aligned} &\|\Psi_1 y_1(t) - \Psi_1 y_2(t)\|_{C_{1-\alpha}}^2 \\ &\leq \frac{\check{N}_{\alpha, \beta}^2 T^{2-2\alpha\beta} K_1}{|(2\alpha-1)(1-2\alpha\beta)|} \|y_1(t) - y_2(t)\|_{C_{1-\alpha}}^2 \\ &\leq \gamma \|y_1(t) - y_2(t)\|_{C_{1-\alpha}}^2 \end{aligned}$$

Where,  $\gamma = \frac{\check{N}_{\alpha, \beta}^2 T^{2-2\alpha\beta} K_1}{|(2\alpha-1)(1-2\alpha\beta)|} < 1$ , hence  $\Psi_1$  is a contraction.

**Lemma (3.7):**

Assume that the assumptions (a) – (e) hold, then the operator  $\Psi_2$  maps bounded sets into bounded sets in  $B_{\check{\delta}}$ .

**Proof:**

Let  $x(t) \in B_{\check{\delta}}$ , for  $t \in [0, T]$ , we have

$$\begin{aligned} &t^{2-2\alpha} E \|\Psi_2 x(t)\|_\beta^2 \\ &\leq 4t^{2-2\alpha} E \left\| \check{T}_\alpha(t)(t)^{\alpha-1} x_0 \right\|_\beta^2 \\ &+ 4t^{2-2\alpha} E \left\| \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} Bu^\theta(s, x) ds \right\|_\beta^2 \\ &+ 4t^{2-2\alpha} E \left\| \int_0^t \left[ \check{T}_\alpha(t-s)(t-s)^{\alpha-1} G\left(s, x(s), \int_0^s g(v, x(v)) dW_{(v)}\right) \right] dW_{(s)} \right\|_\beta^2 \\ &+ 4t^{2-2\alpha} E \left\| \int_0^t \check{T}_\alpha(t-s)(t-s)^{\alpha-1} \sigma(s) dW_{(s)}^H \right\|_\beta^2 \end{aligned}$$

Applying Holder's inequality and by using Ito isometry, Lemmas (2.5), (2.7), (3.3) and the assumptions (a)-(e), we obtain

$$t^{2-2\alpha} E \|\Psi_2 x(t)\|_{\beta}^2 \leq \mathfrak{D}$$

where,  $\mathfrak{D} = 4 \frac{M^2}{(\Gamma(\alpha))^2} E \|x_0\|_{\beta}^2 + 4 \frac{L_B^2 T^{2-2\alpha\beta} N_C \tilde{N}_{\alpha,\beta}^2}{(2\alpha-1)(1-2\alpha\beta)} + 4 \frac{T^{2-2\alpha\beta} \tilde{N}_{\alpha,\beta}^2 D_2}{|(2\alpha-1)(1-2\alpha\beta)|} + 8 \frac{H T^{1+2H-2\alpha\beta}}{|(2\alpha-1)(1-2\alpha\beta)|} \tilde{N}_{\alpha,\beta}^2 C_1$

By taking the supremum over  $t \in [0, T]$  for both sides, we get  $\sup_{t \in [0, T]} t^{2-2\alpha} E \|\Psi_2 x(t)\|_{\beta}^2 \leq \mathfrak{D}$

Therefore, for each  $x \in B_{\delta}$ , we get  $\|\Psi_2 x(t)\|_{C_{1-\alpha}}^2 \leq \mathfrak{D}$ . Then  $\Psi_2$  maps bounded sets into bounded sets in  $B_{\delta}$ .

**Lemma (3.8):**

Assume that the assumptions (a) – (e) hold, then  $\Psi_2$  is a continuous on  $B_{\delta}$ .

**Proof:**

Let  $\{x_n\}_{n=1}^{\infty}$  be a sequence in  $B_{\delta}$  such that  $x_n \rightarrow x$  as  $n \rightarrow \infty$  in  $C_{1-\alpha}([0, T]; L^2(\Omega, X_{\beta}))$ . For each  $t \in [0, T]$ , we have

$$\begin{aligned} & t^{2-2\alpha} E \|(\Psi_2 x_n)(t) - (\Psi_2 x)(t)\|_{\beta}^2 \\ & \leq 2t^{2-2\alpha} E \left\| \int_0^t \left\{ \begin{array}{l} \check{T}_{\alpha}(t-s)(t-s)^{\alpha-1} \\ B[u^{\theta}(s, x_n) - u^{\theta}(s, x)] \end{array} \right\} ds \right\|_{\beta}^2 \\ & + 2t^{2-2\alpha} E \left\| \int_0^t \left\{ \begin{array}{l} \check{T}_{\alpha}(t-s)(t-s)^{\alpha-1} \\ [(\check{G}x_n)(s) - (\check{G}x)(s)] \end{array} \right\} dW(s) \right\|_{\beta}^2 \end{aligned}$$

From Ito isometry and lemma (2.7), we obtain

$$\begin{aligned} & t^{2-2\alpha} E \|(\Psi_2 x_n)(t) - (\Psi_2 x)(t)\|_{\beta}^2 \\ & \leq 2L_B^2 \tilde{N}_{\alpha,\beta}^2 t^{2-2\alpha} \int_0^t \left[ (t-s)^{-2\alpha\beta+2\alpha-2} E \|u^{\theta}(s, x_n) - u^{\theta}(s, x)\|_{\beta}^2 \right] ds \\ & + 2\tilde{N}_{\alpha,\beta}^2 t^{2-2\alpha} \int_0^t \left[ (t-s)^{-2\alpha\beta+2\alpha-2} E \|(\check{G}x_n)(s) - (\check{G}x)(s)\|_{\beta}^2 \right] ds \end{aligned}$$

Therefore, it follows from the continuity of  $G$  and  $u^{\theta}$  that for each  $t \in [0, T]$ ,  $G(s, x_n(s), \int_0^s g(v, x_n(v))dW_{(v)}) \rightarrow G(s, x(s), \int_0^s g(v, x(v))dW_{(v)})$  and  $u^{\theta}(s, x_n) \rightarrow u^{\theta}(s, x)$ , using the Lebesgue dominated convergence theorem that for all  $t \in [0, T]$ , we conclude  $\|(\Psi_2 x_n)(t) - (\Psi_2 x)(t)\|_{C_{1-\alpha}} \rightarrow 0$ , as  $n \rightarrow \infty$ , implying that  $\|\Psi_2 x_n - \Psi_2 x\|_{C_{1-\alpha}} \rightarrow 0$ , as  $n \rightarrow \infty$ . Hence,  $\Psi_2$  is continuous on  $B_{\delta}$ .

**Lemma (3.9):**

If the assumptions (a) – (e) are hold, then for  $x \in B_{\delta}$ , the set  $\{(\Psi_2 x)(t), t \in [0, T]\}$  is equicontinuous.

**Proof:**

Let  $t_1, t_2 \in [0, T]$  such that  $0 < t_1 < t_2 \leq T$ . Then, from the equation (24), we have

$$E \|\Psi_2 x(t_2) - \Psi_2 x(t_1)\|_{\beta}^2 \leq 7E \|[ \check{T}_{\alpha}(t_2) - \check{T}_{\alpha}(t_1) ] x_0\|_{\beta}^2$$

$$\begin{aligned} & + 14E \left\| \int_0^{t_1} (t_2 - s)^{\alpha-1} \begin{bmatrix} \check{T}_{\alpha}(t_2 - s) \\ -\check{T}_{\alpha}(t_1 - s) \end{bmatrix} B u^{\theta}(s, x) ds \right\|_{\beta}^2 \\ & + 14E \left\| \int_0^{t_1} \check{T}_{\alpha}(t_1 - s) \begin{bmatrix} (t_2 - s)^{\alpha-1} \\ -(t_1 - s)^{\alpha-1} \end{bmatrix} B u^{\theta}(s, x) ds \right\|_{\beta}^2 \\ & + 7E \left\| \int_0^{t_2} \check{T}_{\alpha}(t_2 - s) (t_2 - s)^{\alpha-1} B u^{\theta}(s, x) ds \right\|_{\beta}^2 \\ & + 14E \left\| \int_0^{t_1} (t_2 - s)^{\alpha-1} \begin{bmatrix} \check{T}_{\alpha}(t_2 - s) \\ -\check{T}_{\alpha}(t_1 - s) \end{bmatrix} (\check{G}x)(s) dW(s) \right\|_{\beta}^2 \\ & + 14E \left\| \int_0^{t_1} \check{T}_{\alpha}(t_1 - s) \begin{bmatrix} (t_2 - s)^{\alpha-1} \\ -(t_1 - s)^{\alpha-1} \end{bmatrix} (\check{G}x)(s) dW(s) \right\|_{\beta}^2 \\ & + 7E \left\| \int_{t_1}^{t_2} \check{T}_{\alpha}(t_2 - s) (t_2 - s)^{\alpha-1} (\check{G}x)(s) dW(s) \right\|_{\beta}^2 \\ & + 14E \left\| \int_0^{t_1} (t_2 - s)^{\alpha-1} \begin{bmatrix} \check{T}_{\alpha}(t_2 - s) \\ -\check{T}_{\alpha}(t_1 - s) \end{bmatrix} \sigma(s) dW(s) \right\|_{\beta}^2 \\ & + 14E \left\| \int_0^{t_1} \check{T}_{\alpha}(t_1 - s) \begin{bmatrix} (t_2 - s)^{\alpha-1} \\ -(t_1 - s)^{\alpha-1} \end{bmatrix} \sigma(s) dW(s) \right\|_{\beta}^2 \\ & + 7E \left\| \int_{t_1}^{t_2} \check{T}_{\alpha}(t_2 - s) (t_2 - s)^{\alpha-1} \sigma(s) dW(s) \right\|_{\beta}^2 \end{aligned}$$

Now, from Lemma (2.7), noting the fact that for every  $\epsilon > 0$ , there exists  $\tau > 0$  such that, whenever  $t_2 - t_1 < \tau$ , for every  $t_1, t_2 \in [0, T]$ ,  $\|\check{T}_{\alpha}(t_2) - \check{T}_{\alpha}(t_1)\|_{\beta}^2 < \epsilon$ . Therefore, when  $t_2 - t_1 < \tau$ , we have

$$\begin{aligned} & E \|\Psi_2 x(t_2) - \Psi_2 x(t_1)\|_{\beta}^2 \\ & \leq 7\epsilon^2 E \|x_0\|_{\beta}^2 + 14 \frac{L_B^2 \epsilon^2 N_C T^{2\alpha}}{|2\alpha-1|} \\ & + 14 \frac{L_B^2 N_C \tilde{N}_{\alpha,\beta}^2 T^{1-2\alpha\beta}}{|1-2\alpha\beta|} \int_0^{t_1} \left[ \begin{array}{l} (t_2 - s)^{\alpha-1} \\ -(t_1 - s)^{\alpha-1} \end{array} \right]^2 ds \\ & + 7N_C \tilde{N}_{\alpha,\beta}^2 L_B^2 \int_{t_1}^{t_2} (t_2 - s)^{2\alpha-2\alpha\beta-2} ds + 14 \frac{\epsilon^2 D_2 T^{2\alpha}}{|2\alpha-1|} \\ & + 14 \frac{\tilde{N}_{\alpha,\beta}^2 D_2 T^{1-2\alpha\beta}}{|1-2\alpha\beta|} \int_0^{t_1} \left[ (t_2 - s)^{\alpha-1} - (t_1 - s)^{\alpha-1} \right]^2 ds \\ & + 7\tilde{N}_{\alpha,\beta}^2 D_2 \int_{t_1}^{t_2} (t_2 - s)^{2\alpha-2\alpha\beta-2} ds \\ & + 28H\epsilon^2 C_1 \frac{T^{2\alpha+2H-2}}{2\alpha-1} \\ & + 28H T^{2H-1} C_1 \tilde{N}_{\alpha,\beta}^2 \int_0^{t_1} (t_1 - s)^{-2\alpha\beta} \left[ \begin{array}{l} (t_2 - s)^{\alpha-1} \\ -(t_1 - s)^{\alpha-1} \end{array} \right]^2 ds \\ & + 28HT^{2H-1} C_1 \tilde{N}_{\alpha,\beta}^2 \int_{t_1}^{t_2} (t_2 - s)^{2\alpha-2\alpha\beta-2} ds \end{aligned}$$

The right hand of the inequality above tends to 0 as  $t_2 \rightarrow t_1$  and  $\epsilon \rightarrow 0$ . Hence for  $x \in B_{\delta}$ , the set  $\{\Psi_2 x, t \in [0, T]\}$  is equicontinuous.

**Lemma (3.10):**

If the assumption (a) is hold. Then for each  $t \in [0, T]$ , the set  $\tilde{U}(t) = \{\Psi_2 x, x \in B_{\delta}\}$  is relatively compact in  $B_{\delta}$ .

**Proof:**

Let  $t \in (0, T]$  be a fixed and  $0 < \hat{a} < t$ , for every  $\gamma > 0$ ,  $x \in B_{\delta}$ , we define

$$\begin{aligned} \Psi_2^{\hat{a}, \gamma} x(t) &= \int_{\gamma}^{\infty} \alpha r M_{\alpha}(r) \check{T}(t^{\alpha} r)(t)^{\alpha-1} x_0 dr \\ &+ \int_0^{t-\hat{a}} \int_{\gamma}^{\infty} \alpha r M_{\alpha}(r) \check{T}((t-s)^{\alpha} r)(t-s)^{\alpha-1} Bu^{\theta}(t, x) dr ds \\ &+ \int_0^{t-\hat{a}} \int_{\gamma}^{\infty} \alpha r M_{\alpha}(r) \check{T}((t-s)^{\alpha} r)(t-s)^{\alpha-1} (\tilde{G}x)(s) dr dW_{(s)} \\ &+ \int_0^{t-\hat{a}} \int_{\gamma}^{\infty} \alpha r M_{\alpha}(r) \check{T}((t-s)^{\alpha} r)(t-s)^{\alpha-1} \sigma(s) dr dW_{(s)}^H \end{aligned} \tag{25}$$

Then, from the definition of semigroup  $\check{T}(t^{\alpha} r)$ ,  $t^{\alpha} r > 0$ , we can easily be written in the form,  $\check{T}(t^{\alpha} r) = \check{T}(\hat{a}^{\alpha} \gamma)$   $\check{T}(t^{\alpha} r - \hat{a}^{\alpha} \gamma)$ , from the equality (25), we have

$$\begin{aligned} \Psi_2^{\hat{a}, \gamma} x(t) &= \check{T}(\hat{a}^{\alpha} \gamma) \int_{\gamma}^{\infty} \alpha r M_{\alpha}(r) \check{T}(t^{\alpha} r - \hat{a}^{\alpha} \gamma)(t)^{\alpha-1} x_0 dr \\ &+ \check{T}(\hat{a}^{\alpha} \gamma) \int_0^{t-\hat{a}} \int_{\gamma}^{\infty} \left[ \alpha r M_{\alpha}(r) \check{T}((t-s)^{\alpha} r - \hat{a}^{\alpha} \gamma) \right. \\ &\quad \left. (t-s)^{\alpha-1} Bu^{\theta}(t, x) \right] dr ds \\ &+ \check{T}(\hat{a}^{\alpha} \gamma) \int_0^{t-\hat{a}} \int_{\gamma}^{\infty} \left[ \alpha r M_{\alpha}(r) \check{T}((t-s)^{\alpha} r - \hat{a}^{\alpha} \gamma) \right. \\ &\quad \left. (t-s)^{\alpha-1} (\tilde{G}x)(s) \right] dr dW_{(s)} \\ &+ \check{T}(\hat{a}^{\alpha} \gamma) \int_0^{t-\hat{a}} \int_{\gamma}^{\infty} \left[ \alpha r M_{\alpha}(r) \check{T}((t-s)^{\alpha} r - \hat{a}^{\alpha} \gamma) \right. \\ &\quad \left. (t-s)^{\alpha-1} \sigma(s) \right] dr dW_{(s)}^H \end{aligned}$$

Then, from the compactness of  $\check{T}(\hat{a}^{\alpha} \gamma)$ , we obtain that the set  $\{\Psi_2^{\hat{a}, \gamma} x(t), x \in B_{\delta}\}$  is relatively compact in  $X$  for every  $\hat{a}$  such that  $0 < \hat{a} < t$ . Moreover, for  $x \in B_{\delta}$ , we can easily prove that  $\Psi_2^{\hat{a}, \gamma} x(t)$  is convergent to  $\Psi_2 x(t)$  in  $B_{\delta}$ , as  $\hat{a} \rightarrow 0^+$  and  $\gamma \rightarrow 0$ , hence the set  $\tilde{U}(t) = \{\Psi_2 x, x \in B_{\delta}\}$  is relatively compact in  $B_{\delta}$ .

**Lemma (3.11):**

If the assumptions (a) – (e) are hold. Then  $\Psi_2$  is a completely continuous.

**Proof:**

From lemma (3.7) and (3.9), for each  $x \in B_{\delta}$  the operator  $\Psi_2$  is uniformly bounded and the set  $\{\Psi_2 x, t \in [0, T]\}$  is equicontinuous by Applying the Arzela –Ascoli theorem, it results that for  $x \in B_{\delta}$ , the set  $\{\Psi_2 x, t \in [0, T]\}$  is relatively compact.

we obtain that  $\Psi_2$  is a completely continuous.

**Theorem (3.1):**

If the assumptions (a)-(e) are satisfied. Then for each  $\theta > 0$ , the control system (1) has a mild solution on  $[0, T]$ ,

provided that  $\gamma = \frac{\bar{N}_{\alpha, \beta}^2 T^{2-2\alpha\beta} K_1}{|(2\alpha-1)(1-2\alpha\beta)|} < 1$ .

**Proof:**

For any  $\theta > 0$  and for any  $x, y \in B_{\delta}$ ,  $(\Psi_1 y)(t) + (\Psi_2 x)(t) \in B_{\delta}$ , for  $t \in [0, T]$ . By using lemma (3.6) and (3.11) with applying Krasnoselskii’s fixed point theorem, we conclude that the operator  $\Psi_{\theta}$  has a fixed

point, which gives rise to mild solution of system (1) with stochastic control function given in (20). This completes the proof.

**Theorem (3.2):**

If the assumptions (a) – (e) are satisfied. Then the stochastic control system (1) is approximately controllable on  $[0, T]$ .

**Proof:**

For every  $\theta > 0$ , let  $x_{\theta}$  be a fixed point of the operator  $\Psi_{\theta}$  in the space  $C_{1-\alpha}([0, T]; L^2(\Omega, X_{\beta}))$ , which is a mild solution under the stochastic control function in (20) of the stochastic control system (1). Then, we have

$$\begin{aligned} x_{\theta}(T) &= x_T - \epsilon R(\epsilon, \Gamma_0^T) \left( E x_T - \check{T}_{\alpha}(T)(T)^{\alpha-1} x_0 \right) \\ &- \int_0^T \theta R(\theta, \Gamma_s^T) \hat{a}(s) dW_{(s)} \\ &+ \int_0^T \left[ \theta R(\theta, \Gamma_s^T) \check{T}_{\alpha}(T-s)(T-s)^{\alpha-1} \right. \\ &\quad \left. F\left(s, x_{\theta}(s), \int_0^s h(s, r, x_{\theta}(r)) dr \right) \right] ds \\ &+ \int_0^T \left[ \theta R(\theta, \Gamma_s^T) \check{T}_{\alpha}(T-s)(T-s)^{\alpha-1} \right. \\ &\quad \left. G\left(s, x_{\theta}(s), \int_0^s g(v, x_{\theta}(v)) dW_{(v)} \right) \right] dW_{(s)} \\ &+ \int_0^T \theta R(\theta, \Gamma_s^T) \check{T}_{\alpha}(T-s)(T-s)^{\alpha-1} \sigma(s) dW_{(s)}^H \end{aligned} \tag{25}$$

By using the assumptions (c) the function  $F$  is uniformly bounded, such that  $E \left\| F(t, x, \int_0^t h(t, s, x) ds) \right\|^2 \leq D_1$ , for any  $0 \leq s \leq T$ . Then, for all  $(s, \omega) \in [0, T] \times \Omega$ , there is a subsequence of sequence  $\{F(s, x_{\theta}(s), \int_0^s h(s, r, x_{\theta}(r)) dr)\}$  denoted by  $\{F(s, x_{\theta}(s), \int_0^s h(s, r, x_{\theta}(r)) dr)\}$  which is weakly converging to say  $F(s)$  in  $X$ . Also, there is a subsequence of sequence  $\{G(s, x_{\theta}(s), \int_0^s g(v, x_{\theta}(v)) dW_{(v)})\}$  denoted by  $\{G(s, x_{\theta}(s), \int_0^s g(v, x_{\theta}(v)) dW_{(v)})\}$  which is weakly converging to say  $G(s)$  in  $L_2(K, X)$ . On the other hand, from the assumption (b), for all  $0 \leq s < T$  the operator  $\theta R(\theta, \Gamma_s^T) \rightarrow 0$  Strongly as  $\theta \rightarrow 0^+$  and  $\|\theta R(\theta, \Gamma_s^T)\| \leq 1$ .

$$\begin{aligned} &E \|x_{\theta}(T) - x_T\|^2 \\ &\leq 8E \left\| \theta R(\theta, \Gamma_s^T) \left( E x_T - \check{T}_{\alpha}(T)(T)^{\alpha-1} x_0 \right) \right\|^2 \\ &+ 8E \left\| \int_0^T \theta R(\theta, \Gamma_s^T) \phi(s) dW_{(s)} \right\|^2 \\ &+ 8E \left\| \int_0^T R(\theta, \Gamma_s^T) \hat{H}(s) dW_{(s)}^H \right\|^2 \\ &+ 8E \left\| \int_0^T \left[ R(\theta, \Gamma_s^T) \check{T}_{\alpha}(T-s)(T-s)^{\alpha-1} \right. \right. \\ &\quad \left. \left. \left[ F\left(s, x_{\theta}(s), \int_0^s h(s, r, x_{\theta}(r)) dr \right) - F(s) \right] \right] ds \right\|^2 \\ &+ 8E \left\| \int_0^T \theta R(\theta, \Gamma_s^T) \check{T}_{\alpha}(T-s)(T-s)^{\alpha-1} F(s) ds \right\|^2 \end{aligned}$$



$$+8E \left\| \int_0^T \left[ \begin{array}{c} \theta R(\theta, \Gamma_s^T) \check{T}_\alpha (T-s)(T-s)^{\alpha-1} \\ G \left( s, x_\theta(s), \int_0^s g(v, x_\theta(v)) dW_{(v)} \right) \\ -G(s) \end{array} \right] dW_{(s)} \right\|^2$$

By using the Lebesgue dominated convergence theorem, we obtain  $E\|x_\theta(T) - x_T\|^2 \rightarrow 0$  as  $\theta \rightarrow 0^+$ . This gives the approximate controllability.

### References

[1] Abid S. H., Hasan S. Q. and Quaez U. J. "Approximate controllability of Fractional Stochastic Integro-Differential Equations Driven by Mixed Fractional Brownian Motion", American Journal of Mathematics and Statistics 2015, Vo. 2, PP:72-81, 2015.

[2] Ahmed M. Hamdy, "Approximate Controllability Of Impulsive Neutral Stochastic Differential Equations With Fractional Brownian Motion in A Hilbert Space", Advances in Difference Equations, Springer Open Journal, 2014:113, 2014.

[3] Balachandran K., Kiruthika S. and Trujillo J. "On Fractional Impulsive Equations Of Sobolev Type with Nonlocal Conditions in Banach Spaces", Computers and Mathematics with Applications, No. 62, PP: 1157-1165, 2011.

[4] Engel K. J. and Nagel R. "One Parameter Semigroup For Linear Evolution Equations", Springer-Verlag, New York, Berlin, 2000.

[5] Gani J., Heyde C.C., Jagers P. and Kurtz T.G., "Probability and Its Applications", Springer-Verlag London Limited, 2008.

[6] Gripenberg, G. and Norros I., "On The Prediction Of Fractional Brownian Motion", Journal of Applied Probability, Vol. 33, No. 2, PP: 400-410, 1996.

[7] Guendouzi T. and Idrissi S., "Approximate Controllability of Fractional Stochastic Functional Evolution Equations Driven By A Fractional Brownian Motion", Romai J., Vo.8, No.2, PP:103-117, 2012.

[8] Kerboua M., Debbouche A. and Baleanu D., "Approximate Controllability Of Sobolev Type Fractional Stochastic Nonlocal Nonlinear Differential Equations in Hilbert Spaces", Electronic Journal Of Qualitative Theory of Differential Equations, No. 58, PP: 1-16, 2014.

[9] Li C., Qian D. and Chen Y., "On Riemann- Liouville and Caputo Derivatives", Hindawi Publishing Corporation, Vol. 2011, Article ID 562494, 15 pages.2011.

[10] Li K., "Stochastic Delay Fractional Evolution Equations Driven By Fractional Brownian Motion", Mathematical Method in The Applied Sciences, 2014.

[11] Madsen Henrik, "ito integrals", 2006.

[12] Mahmudov N. and Zorlu S., "Approximate Controllability Of Fractional Integro-Differential Equations Involving Nonlocal Initial Conditions", Boundary Value Problems, Springer Open Journal, 2013:118, 2013.

[13] Mahmudov N., "Controllability of Linear Stochastic Systems in Hilbert Spaces", Journal of Mathematical Analysis and Applications Vo. 259, PP: 64-82, 2001.

[14] Mishura Y. S., "Stochastic Calculus for Fractional Brownian Motion and Related Processes", Lect.Notes in Math., 1929, Springer, 2008.

[15] Nourdin I., "Select Aspects of fractional Brownian Motion", Springer-Verlag Italia, 2012.

[16] Nualart D., "Fractional Brownian motion: stochastic calculus and Applications", Proceedings of the International Congress of Mathematicians, Madrid, Spain, European Mathematical Society, 2006.

[17] Pazy, A., "Semigroup of Linear Operator and Applications to Partial Differential Equations", Springer-Verlag, New York, 1983.

[18] Podlubny I., "Fractional Differential Equations", Academic Press, San Diego. California, USA, 1999.

[19] Sakthivel R., "Approximate Controllability Of Impulsive Stochastic Evolution Equations", Funkcialaj Ekvacioj, Vol. 52(2009), PP:381-393, 2009.

[20] Zhou Y., Wang J. and Feckan M. "Controllability Of Sobolev Type Fractional Evolution systems ", Dynamics of PDE, Vol. 11, No. 1, PP: 71-87, 2014.